



BIOMAX PLANT ACTIVATOR

SUMMARY OF ANALYSIS

A talc-based powder mixture of 1% W.P. of *Trichoderma harzianum* and *Trichordema viridae* (1 x 10¹¹ cfu/g)

APPLICATION RATES

Soil treatment Apply 2kg per hectare or 20g per 100m² along with fertiliser for first fertigation. For further application, mix 2kg per hectare or 20g per 100m² in water and spray every 2-4 weeks depending on crop growth cycle.

Seeds treatment

Mix 6-10g with water to make a liquid mixture.

Moisten 1kg of seeds with the liquid mixture for 15-30 minutes before sowing.

Nursery treatment

Mix 5-10g with 1kg of germination medium. Transfer the germination medium into germination tray. Apply 10-20g per 100m² of nursery bed before planting. Acclimatize seedlings in a mixture of 5-10g per 1kg of nursery mix.

Seedling treatment

Mix 10gm in 1 litre of water and dip the seedlings in the solution for 10 minutes before planting.

Side dressing

Mix 2kg with water and spray the mixed solution to 1 hectare of planting land.

Please contact us for any other application.

Do not use in dry soil or media. Do not expose product to strong sunlight. Avoid using with chemical fungicides. Use the treated soil within 1 week. Use overnight tap water for a better result.

